


From Diagnosis to Daily Functioning: Translating ADHD Data into Action



Sam Goldstein, Ph.D.

Assistant Clinical Professor
University of Utah School of Medicine



1

Goals for This Presentation

- . Interpret ADHD assessment findings through a functional lens.
- . Translate diagnostic data into individualized intervention plans.
- . Identify strengths and challenges that affect daily performance.
- . Match assessment findings to evidence-based interventions.
- . Measure success using functional outcomes rather than symptom reduction alone.

2

An ADHD evaluation should do more than establish a diagnosis. It should provide a roadmap to improve daily functioning.

- Today's presentation focuses on one of the greatest challenges in ADHD assessment: translating clinical findings into practical recommendations that improve everyday functioning.
- Too often, evaluations end with a diagnosis and a list of test scores. Families leave with valuable information but little guidance about what those results actually mean in everyday life.
- Throughout this webinar, we'll shift our perspective. Instead of asking, "Does this person have ADHD?" we'll ask, "How does ADHD affect this person's daily functioning, and what specific interventions can improve outcomes?"
- Our goal is to bridge the gap between assessment and intervention by showing how evaluation data can guide individualized supports across home, school, work, and social settings.
- Every assessment should answer the question:
"What should this individual do differently tomorrow because of what we learned today?"

3

Today's objectives emphasize application over diagnosis. Most clinicians are comfortable determining whether diagnostic criteria have been met. The greater challenge is translating assessment findings into practical recommendations that improve daily life.

4

Assessment should always answer two questions:

What explains the problem?
What should we do about it?

5

Why We Evaluate ADHD

- Families rarely seek evaluations because of DSM criteria.
- The real referral question is: "How can life work better?"
- They seek evaluations because:
 - Homework takes hours.
 - Morning routines are chaotic.
 - Relationships are strained.
 - Work performance is inconsistent.
 - Everyday life feels overwhelming.

6

A comprehensive evaluation should therefore focus on explaining these functional difficulties and identifying practical strategies to reduce impairment. The diagnosis provides context. Intervention changes lives. Symptoms bring people into the office. Functional impairment determines treatment.

7

Beyond the Diagnosis

- A diagnosis answers:
 - Does this individual meet the criteria for ADHD?
- An evaluation should also answer:
 - Why are these difficulties occurring?
 - Where do they interfere most?
 - Which strengths can be leveraged?
 - Which supports are most likely to help?
 - How will progress be measured?

8

Diagnosis explains the problem.

Intervention improves functioning.

9

ADHD as a Disorder of Self-Regulation Involving 3 Key Challenges

- More experiential repetition is required to make behavior automatic.
- More stimulation or interest is required to come to and stick with a non-desired task.
- Repetition quickly causes satiation.

10

Modern conceptualizations emphasize ADHD as a disorder of executive self-regulation.

Executive functions include:

- Response inhibition
- Working memory
- Planning
- Organization
- Cognitive flexibility
- Emotional regulation
- Self-monitoring
- Goal-directed persistence

11

While attention difficulties remain central to ADHD, research increasingly highlights impairments in executive functioning and self-regulation.

These skills enable individuals to manage behavior to achieve long-term goals.

Viewing ADHD through an executive-function framework helps explain why individuals may understand what needs to be done but struggle to carry it out consistently.

12

Knowing is not the same as doing.

ADHD primarily affects performance, not intelligence.

13

Executive Function and Everyday Performance

- Executive function weaknesses affect:
 - Starting tasks
 - Sustaining effort
 - Organizing information
 - Managing time
 - Regulating emotions
 - Monitoring performance
 - Completing long-term projects

Executive function problems are often the strongest predictor of everyday impairment.

14

Executive functioning influences nearly every aspect of daily life. Difficulties initiating tasks, estimating time, organizing materials, and regulating emotions contribute to the functional impairments associated with ADHD.

Understanding these processes enables clinicians to recommend supports that compensate for specific executive weaknesses rather than relying on generic advice.

Executive function problems are often the strongest predictor of everyday impairment.

15

Components of a Comprehensive ADHD Evaluation

A high-quality evaluation integrates multiple sources of information:

- Clinical interview
- Developmental history
- Behavioral observations
- Rating scales
- Cognitive assessment
- Academic testing (when indicated)
- Functional assessment
- Collateral information

No single measure diagnoses ADHD. Comprehensive evaluations synthesize information from multiple sources to understand the individual's functioning across settings.

This multimethod approach increases diagnostic accuracy and provides a richer understanding of strengths, challenges, and contextual factors influencing performance.

The goal is not simply confirming ADHD but identifying the factors that contribute to functional impairment.

16

Two individuals with identical ADHD symptom counts may function very differently depending on cognitive strengths, family support, educational environment, and co-occurring conditions.

Effective intervention planning requires understanding the whole person rather than focusing solely on symptom severity.

This broader perspective leads to more individualized and successful recommendations.

Treat the person, not the diagnosis.

17

Assessment should identify:

- . Strengths
- . Weaknesses
- . Environmental demands
- . Protective factors
- . Risk factors
- . Comorbid conditions
- . Available supports
- . Areas of greatest impairment

18

Every recommendation should answer four questions:

- What is the problem?
- Where does it occur?
- What intervention addresses it?
- How will we know it worked?

This four-step framework serves as the foundation for the remainder of the presentation. Every assessment finding should be linked to a functional consequence, an evidence-based intervention, and a measurable outcome. This approach ensures that evaluation results lead directly to meaningful changes in daily life and provides a clear method for monitoring progress over time.

19

The Clinical Interview: The Foundation of Every Evaluation

ADHD is diagnosed by history, not by testing alone.

The clinical interview provides context for every test score.

While standardized assessments are valuable, they cannot replace a thorough understanding of the individual's developmental course and current functioning.

Interview multiple informants whenever possible, as parents, teachers, spouses, and employers often provide complementary perspectives.

Look beyond symptoms to identify patterns of success and failure, environmental demands, coping strategies, and previous interventions. Ask not only what is difficult but also when the individual performs well.

20

Rating scales quantify observed behavior but should never be interpreted in isolation.

Differences between home and school ratings often provide valuable clinical information. For example, significant symptoms at school but not at home may suggest environmental influences or varying task demands.

Discuss discrepancies openly rather than viewing them as contradictions. ADHD symptoms fluctuate depending on structure, motivation, fatigue, and expectations.

Differences between raters are often clinically meaningful, not errors.

21

Using Rating Scales

- Rating scales are an essential component of a comprehensive neuropsychological evaluation.
- Although they do not establish a diagnosis independently, they provide valuable information about how symptoms present across environments, their severity, and their functional impact.

22

Compare behaviors across settings

- Determine whether symptoms are present at home, school, and in the community.
- Identify discrepancies between parent, teacher, self-, and clinician reports.
- Evaluate the consistency of concerns across multiple informants to strengthen diagnostic confidence.
- Differentiate situational problems from pervasive neurodevelopmental or psychiatric disorders.

23

Assess symptom severity

- Quantify the frequency and intensity of cognitive, behavioral, emotional, and social symptoms.
- Compare an individual's functioning with age-based normative data. Identify clinically significant areas requiring intervention.
- Establish a baseline for future comparison.

24

Evaluate executive functioning

- Assess everyday difficulties with attention, working memory, inhibition, cognitive flexibility, planning, organization, initiation, emotional regulation, and self-monitoring.
- Compare observed test performance with real-world functioning. Identify strengths and weaknesses that may not emerge during structured testing.
- Guide recommendations for school, home, and therapeutic interventions.

25

Monitor treatment outcomes

- Measure changes following behavioral, educational, medical, or psychological interventions.
- Evaluate the effectiveness of medication and other treatment strategies over time.
- Document progress objectively using repeated standardized measures.
- Assist in modifying treatment plans based on response to intervention.

26

Support differential diagnosis

- Identify symptom patterns that distinguish ADHD, anxiety, depression, autism spectrum disorder, sleep disorders, traumatic brain injury, and other neurodevelopmental conditions.
- Recognize when symptoms may reflect emotional distress, environmental factors, or medical conditions rather than a primary neurocognitive disorder.

27

Assess functional impairment

- Determine how symptoms affect academic achievement, social relationships, adaptive functioning, and daily living skills.
- Provide objective evidence supporting eligibility for educational accommodations and clinical services.

28

Facilitate communication

- Provide a common language for discussing concerns with families, educators, physicians, and other professionals.
- Enhance interdisciplinary collaboration by supplying standardized, interpretable data.

29

Guide clinical decision-making

- Integrate rating scale findings with interview, history, behavioral observations, cognitive testing, and medical information.
- Improve diagnostic accuracy and inform evidence-based recommendations.

30

Behavioral observations help bridge test performance and everyday functioning. A child who frequently loses focus, rushes through tasks, or gives up easily during testing may demonstrate similar behaviors in school and at home.

Likewise, adults who repeatedly interrupt, lose track of conversations, or become overwhelmed by complex tasks often reveal executive function challenges that standardized scores alone may not capture.

Describe behaviors objectively and relate them directly to likely functional consequences.

The way someone completes testing is often as informative as the score itself.

31

Integrating Multiple Rating Scales into Assessment

Each instrument contributes a distinct perspective to the evaluation process:

Instrument	Primary Clinical Question
Conners-4	<i>Are symptoms consistent with ADHD or related problems in attention, behavior regulation, and executive functioning?</i>
CEFI	<i>How effectively does the individual manage executive functioning demands in everyday life?</i>
RSI	<i>To what extent do the identified symptoms interfere with academic, social, family, adaptive, or occupational functioning?</i>
ASRS	<i>Are there clinically significant characteristics of autism involving social communication, restricted interests, behavioral rigidity, and sensory processing?</i>
MASC-2	<i>Are anxiety symptoms contributing to emotional, behavioral, social, or academic difficulties?</i>
CDI-2	<i>Are depressive symptoms affecting mood, motivation, cognition, interpersonal relationships, or overall functioning?</i>

32

Clinical Purpose	Conners-4	CEFI	Rating Scale of Impairment (RSI)	ASRS (Autism Spectrum Rating Scales)	MASC-2	CDI-2
Primary Purpose	Assess ADHD symptoms and related behavioral, emotional, and learning concerns.	Assess executive functioning in everyday settings.	Measure the degree to which symptoms interfere with daily functioning.	Assess behaviors associated with Autism Spectrum Disorder, including social communication and restricted/repetitive behaviors.	Assess anxiety symptoms across multiple domains.	Assess depressive symptoms and associated impairment.
Compare Behaviors Across Settings	✓ Parent, Teacher, and Self forms compare behavior across home, school, and community.	✓ Parent, Teacher, and Self forms compare executive functioning across environments.	✓ Parent, Teacher, and Self ratings compare functional impairment across settings.	✓ Parent and Teacher forms compare autism-related behaviors across settings. Adult version includes self- and informant-report.	✓ Parent and Self forms compare anxiety symptoms across settings.	✓ Parent and Self forms compare depressive symptoms across settings.
Assess Symptom Severity	✓ Standardized scores for inattention, hyperactivity/impulsivity, executive dysfunction, aggression, peer relations, and learning problems.	✓ Quantifies severity of executive functioning strengths and weaknesses.	✓ Quantifies severity of functional impairment resulting from symptoms.	✓ Measures severity of autism-related social communication deficits, behavioral rigidity, sensory sensitivity, and unusual behaviors.	✓ Measures overall anxiety and specific anxiety domains.	✓ Measures severity of depressive symptoms including mood, self-esteem, interpersonal difficulties, and negative affect.

33

Clinical Purpose	Conners-4	CEFI	Rating Scale of Impairment (RSI)	ASRS (Autism Spectrum Rating Scales)	MASC-2	CDI-2
Evaluate Executive Functioning	✓ Includes Executive Function scales assessing planning, organization, inhibition, working memory, and self-regulation.	✓ Comprehensive assessment of attention, planning, organization, working memory, flexibility, inhibition, emotional regulation, initiation, and self-monitoring.	✓ Evaluates the functional impact of executive weaknesses.	Indirectly identifies executive challenges commonly associated with ASD (e.g., flexibility, organization, self-regulation).	Indirectly reflects executive inefficiency related to anxiety.	Indirectly reflects executive slowing and motivational deficits associated with depression.
Assess Functional Impairment	✓ Measures impact on academics, family functioning, peer relationships, and behavior.	✓ Assesses impact of executive functioning on success in home, school, and community settings.	✓ Primary purpose: Measures impairment in academic, family, social, occupational, and adaptive functioning regardless of diagnosis.	✓ Identifies functional impact of social communication deficits and behavioral rigidity on everyday functioning.	✓ Evaluates impairment associated with anxiety in school, family, and social settings.	✓ Evaluates impairment associated with depressive symptoms in daily functioning.
Support Differential Diagnosis	✓ Helps differentiate ADHD from learning, behavioral, emotional, and developmental disorders when interpreted with other data.	✓ Helps distinguish executive dysfunction associated with ADHD, ASD, learning disorders, acquired brain injury, emotional disorders, and neurological conditions.	✓ Clarifies whether symptoms produce clinically meaningful impairment necessary for diagnosis and intervention planning.	✓ Assists in distinguishing ASD from ADHD, language disorders, anxiety disorders, social communication disorder, intellectual disability, and emotional disorders.	✓ Identifies anxiety disorders that may mimic or coexist with ADHD, ASD, learning disorders, or depression.	✓ Identifies depressive disorders that may contribute to poor concentration, fatigue, irritability, withdrawal, or academic decline.

34

Clinical Purpose	Conners-4	CEFI	Rating Scale of Impairment (RSI)	ASRS (Autism Spectrum Rating Scales)	MASC-2	CDI-2
Monitor Treatment Outcomes	Tracks medication & behavioral response	Tracks EF change	Tracks functional improvement	Tracks social/adaptive gains	Tracks anxiety response	Tracks depression response
Guide Educational Planning	Identifies classroom supports	Guides EF accommodations	Documents impairment for services	Identifies ASD-related supports	Guides anxiety accommodations	Guides mood-related supports
Facilitate Communication	Standardizes parent, school & provider reports	Common EF framework	Documents functional impact	Standardizes ASD findings	Standardizes anxiety findings	Standardizes depression findings
Major Strength	Broad ADHD & behavioral profile	Comprehensive real-world EF	Measures functional impairment	Comprehensive ASD profile	Broad anxiety assessment	Comprehensive depression assessment

35

Clinical Integration

- Compare symptom presentation across **multiple informants and settings**.
- Quantify the **severity** of behavioral, emotional, executive, and developmental concerns.
- Evaluate **real-world executive functioning** beyond performance-based cognitive testing.
- Determine whether symptoms result in **clinically significant impairment**, a key requirement for diagnosis.
- Differentiate among ADHD, autism spectrum disorder, anxiety disorders, depressive disorders, learning disorders, and other neurodevelopmental or psychiatric conditions.
- Identify co-occurring conditions that may complicate diagnosis and treatment. Establish objective **baseline measures** for monitoring treatment outcomes.
- Guide recommendations for school accommodations, behavioral interventions, psychotherapy, medication management, and referrals.
- Facilitate communication among families, educators, physicians, psychologists, therapists, and other members of the multidisciplinary team.

36

Cognitive testing is not used to diagnose ADHD directly. Instead, it helps explain how the individual processes information and why specific functional problems occur.

Rather than emphasizing IQ scores, focus on meaningful patterns. A significant weakness in working memory or processing speed may explain academic struggles, incomplete work, or difficulty following complex instructions despite average or above-average reasoning abilities.

Families benefit most when findings are translated into practical implications.

Test scores explain *how* someone learns—not whether they can succeed.

37

Working memory refers to the ability to temporarily hold and manipulate information while completing a task.

It affects virtually every daily activity, from remembering homework assignments to following multi-step workplace instructions.

Parents often interpret these behaviors as laziness or carelessness when they are actually manifestations of executive dysfunction.

External supports compensate for internal working memory weaknesses.

Don't expect individuals with weak working memory to "remember better." Help them remember differently.

38

Processing speed reflects efficiency rather than intelligence. Many individuals with ADHD understand material perfectly but require additional time to organize thoughts, respond, or complete written work.

Avoid interpreting slow work as poor motivation. Instead, identify environmental adjustments that allow the individual to demonstrate what they know.

Measure competence separately from speed.

39

Executive functioning should be conceptualized as a profile rather than a single ability.

Two individuals with ADHD may have very different patterns of strengths and weaknesses.

One person may primarily struggle with organization while another experiences significant emotional impulsivity.

Tailoring interventions to the individual's executive profile increases treatment effectiveness.

The intervention should match the executive weakness—not simply the diagnosis.

40

Families often leave evaluations focused entirely on deficits. This can unintentionally reinforce hopelessness and lower self-esteem. Identifying strengths provides opportunities for success and increases engagement in intervention.

A student with exceptional verbal abilities may benefit from oral presentations rather than lengthy written assignments. A highly creative adult may thrive in flexible work environments emphasizing innovation.

Strengths should actively shape treatment recommendations.

Interventions succeed faster when built on existing strengths.

41

The final goal of assessment is a functional formulation, a concise explanation of how cognitive, behavioral, and environmental factors interact to produce everyday challenges. This formulation becomes the bridge between diagnosis and intervention.

When recommendations clearly flow from the formulation, families, educators, and other professionals are far more likely to implement them consistently.

A good report doesn't simply describe difficulties—it explains them and points toward practical solutions.

42

Instead of writing:

"Weak executive functioning."

Write:

"Executive functioning weaknesses interfere with independent organization, task completion, and time management at school and home. These difficulties contribute to missed assignments and increased family conflict."

Functional formulation answers:

What?

Why?

Where?

What next?

43

The DSM requires evidence of impairment because symptoms alone are insufficient. A person may meet symptom criteria yet function relatively well, while another with fewer symptoms experiences significant disruption.

Intervention should therefore target the areas where ADHD interferes with meaningful daily activities.

Ask, "Where is this person unable to meet everyday expectations?" Those are the areas that deserve intervention.

Treat impairment, not test scores.

44

Every evaluation should identify strengths.

Examples include:

- Creativity
- Verbal reasoning
- Curiosity
- Humor
- Problem-solving
- Persistence
- Innovation
- Empathy
- Strengths become intervention tools.

45

Families often leave evaluations that focus entirely on deficits. This can unintentionally reinforce hopelessness and lower self-esteem. Identifying strengths creates opportunities for success and increases engagement in intervention.

A student with exceptional verbal abilities may benefit from oral presentations rather than lengthy written assignments. A highly creative adult may thrive in flexible work environments that emphasize innovation.

46

Diagnosis describes the disorder. Function describes its impact.

The DSM requires evidence of impairment because symptoms alone are insufficient. A person may meet symptom criteria yet function relatively well, while another with fewer symptoms experiences significant disruption.

Intervention should therefore target the areas where ADHD interferes with meaningful daily activities.

Ask, "Where is this person unable to meet everyday expectations?" Those are the areas that deserve intervention.

Treat impairment not test scores.

47

Home Functioning: Where ADHD Is Often Most Visible

- Families frequently describe the home as the most stressful environment because executive demands are constant.
- Children are expected to remember routines, manage time, regulate emotions, and complete responsibilities with less external structure than they receive at school.
- The goal is to redesign the environment so success becomes more likely.
- Replace repeated verbal reminders with visual routines whenever possible.

48

School Functioning: Beyond Grades

- Grades often underestimate the impact of ADHD.
- Many students achieve average grades through extraordinary effort while experiencing high levels of stress, conflict, and exhaustion.
- Assessment should determine how much support is required for success, not simply whether success occurs.
- Good grades do not always indicate good functioning.

49

Workplace Functioning in Adults

- Adult ADHD often becomes apparent when workplace demands exceed executive capacity.
- Difficulty managing multiple projects, organizing information, or estimating time can affect career advancement despite strong knowledge and motivation.
- Recommendations should focus on modifying workflows, using external organizational systems, and reducing unnecessary executive demands.
- Productivity often improves when systems change, not when motivation increases.

50

Social and Relationship Functioning

- Many individuals with ADHD struggle socially because executive weaknesses influence communication, emotional regulation, and self-monitoring.
- Interrupting conversations, forgetting commitments, or reacting impulsively can strain relationships.
- Intervention should include social coaching, emotional regulation skills, and improved communication strategies.
- Relationship problems are often executive function problems expressed socially.

51

Emotional Regulation: An Often Overlooked Dimension

- Although emotional dysregulation is not a DSM diagnostic criterion, it is one of the most impairing aspects of ADHD for many individuals.
- Emotional reactions are often immediate, intense, and difficult to control, leading to conflict at home, school, and work.
- Teaching self-regulation strategies should be part of every comprehensive treatment plan.
- Many ADHD-related crises begin as regulation problems rather than attention problems.

52

Sleep, Health, and Lifestyle Factors

- Lifestyle factors significantly influence ADHD symptoms.
- Poor sleep, inconsistent routines, and unhealthy lifestyle habits can magnify executive functioning difficulties.
- Before assuming worsening ADHD, clinicians should examine these contributing factors.
- Behavioral interventions often become more effective after sleep and health issues are addressed.
- Never ignore sleep when evaluating attention.

53

Identifying Strengths That Promote Success

How can these strengths support intervention?

- Strengths are not simply encouraging observations, they are clinical tools. They help increase engagement, improve motivation, and create more effective interventions.
- A student who enjoys technology may benefit from digital planning tools. An adult with excellent verbal skills may use voice-to-text software instead of relying on written notes.
- Successful intervention amplifies strengths while supporting weaknesses.

54

Setting Functional Goals

Goals should be:

- Specific
- Measurable
- Relevant
- Functional
- Time-limited

Instead of:

- "Improve attention."

Use:

- "Complete homework independently four nights each week."
- Families frequently identify broad goals that are difficult to measure. Functional goals focus on observable behaviors that matter in daily life.
- Well-defined goals also make progress easier to monitor and allow clinicians to adjust interventions based on objective outcomes.
- If progress cannot be measured, it cannot be managed.

55

From Assessment to Individualized Treatment Planning

Every treatment plan should answer four questions:

1. What is the functional problem?
2. What assessment finding explains it?
3. What intervention addresses it?
4. How will success be measured?

Assessment → Understanding → Intervention → Outcome

56

This framework summarizes the purpose of a comprehensive ADHD assessment. Rather than producing generic recommendations, clinicians should directly connect evaluation findings to individualized intervention strategies.

For example, if working memory weaknesses contribute to forgotten assignments, the intervention should emphasize external memory supports, and success should be measured through improved homework completion.

This logical progression transforms assessment into meaningful clinical action.

Every recommendation should be traceable to a specific assessment finding.

57

Building an Individualized Treatment Plan

Effective treatment plans are:

- . Individualized
- . Strengths-based
- . Function-focused
- . Evidence-informed
- . Collaborative
- . Flexible over time

Treatment should address both symptoms and everyday functioning.

58

No two individuals with ADHD have identical needs. Even when symptom presentations are similar, differences in cognitive profile, environmental demands, motivation, family resources, and co-occurring conditions require individualized intervention planning.

A comprehensive treatment plan integrates assessment findings, patient goals, family priorities, and evidence-based interventions into one coordinated roadmap.

Remember that the objective is not simply symptom reduction, it is improving participation in meaningful life activities.

Successful intervention amplifies strengths while supporting weaknesses.

59

Medication

- Medication remains one of the most effective treatments for core ADHD symptoms, but it should not be viewed as a stand-alone intervention.
- Improved attention does not automatically create better organizational skills, healthier relationships, or stronger study habits.
- Clinicians should explain that medication increases the opportunity for success while behavioral strategies teach individuals how to capitalize on that opportunity.
- Medication opens the door—skills and supports help individuals walk through it.
- Medication improves performance capacity; it does not replace skill development.

60

Behavioral Interventions

- Behavioral interventions are particularly effective because ADHD affects self-regulation rather than knowledge.
- Most individuals already know what they should do; they struggle with consistently doing it.
- Providing immediate reinforcement, reducing delays between behavior and consequences, and increasing environmental cues improve follow-through and reduce frustration.
- Behavioral interventions should be proactive rather than reactive.
- Reward the behaviors you want to see more often.

61

Parent Training

- Parent training is among the strongest evidence-based interventions for children with ADHD.
- Rather than attempting to eliminate ADHD behaviors, parents learn strategies that reduce conflict, increase cooperation, and build executive functioning through structured support.
- Encourage parents to use brief, specific instructions, visual schedules, and positive feedback more often than correction.
- Reducing family stress benefits everyone.
- Structure reduces conflict more effectively than repeated reminders.

62

School

- School accommodations should never be generic. Instead, they should directly address the functional challenges identified during assessment.
- For example, a student with weak working memory benefits from written instructions, while one with processing speed weaknesses may require extended time.
- Organizational coaching may be more effective than reducing workload for students whose primary challenge is planning.
- Link every accommodation to a documented need.
- An accommodation should solve a specific problem—not simply be available because of a diagnosis.

63

ADHD Coaching

- Executive function coaching complements therapy by focusing on practical implementation.
- Coaches help individuals break large goals into manageable steps, establish routines, monitor progress, and develop sustainable organizational habits.
- Coaching works particularly well when linked directly to assessment findings and real-world goals.
- Rather than giving advice, coaches teach systems that individuals can continue using independently.
- Teach strategies, not dependence.

64

Every evaluation should answer:

- . What explains the person's difficulties?
- . What strengths can be leveraged?
- . What interventions are most likely to improve functioning?
- . How will we know the plan is working?

65

The purpose of assessment is not to describe a person, it is to improve that person's everyday life. Assessment should change lives.

66

Remember:

Assessment is the beginning, not the end.
Function matters more than symptoms alone.
Strengths should guide intervention.
Recommendations must be individualized.
Progress should be measured objectively.
Collaboration improves outcomes.

These six principles summarize my entire presentation.
Regardless of the assessment tools used, clinicians should
always connect findings to functional implications, evidence-
based recommendations, and measurable outcomes.

67

Every evaluation should leave families with a clear understanding of:

- Why problems occur
- What can be done
- Who will implement the plan
- How progress will be measured

Information becomes valuable only when it changes practice.

68

